

WHAT'S NEW IN INTENSIVE CARE



Ten key points about ICU palliative care

Jeffrey D. Edwards^{1*}, Louis P. Voigt² and Judith E. Nelson^{2,3}

© 2016 Springer-Verlag Berlin Heidelberg and ESICM

Introduction

Palliative care is a core component of comprehensive care for patients facing critical illness, regardless of age, diagnosis, or prognosis. Key domains of intensive care unit (ICU) palliative care include relief of distressing symptoms, effective communication about care goals, patient-focused decision-making, caregiver support, and continuity across care settings. In this article, we highlight ten evidence-based principles of palliative care that help support optimal critical care practice.

1. Palliative care is relevant for all critically ill patients

Although early efforts in ICU palliative care focused on improving the quality of dying and death, the scope of this care extends more broadly to encompass needs of patients pursuing full critical care/life-supporting therapies and expected to survive. Palliative care addresses issues across the spectrum of critical illness within and beyond the ICU, including symptom distress (physical, psychological, spiritual), exchange of information for shared decision-making, and burdens for caregivers.

2. Palliative care is the responsibility of all clinicians in the ICU

“Primary/generalist” palliative care is the responsibility of all ICU clinicians, who require basic knowledge and skills for symptom management, clear and sensitive communication, and shared decision-making based on patients’ values, goals, and preferences [1]. Although palliative care specialists are increasing in number in some countries and can provide essential input in the ICU where available, the workforce of such specialists will remain inadequate to meet increasing needs. A mixed model, in

which primary palliative care is combined with specialist palliative care contributions, is recommended.

3. ICU clinicians can and are obliged to ensure patient comfort at the end of life

As most ICU deaths follow a decision to limit life support, critical care clinicians require competence in the applied and ethical aspects of withholding/withdrawing intensive care interventions [2, 3]. These include the use of sedatives, analgesics, and nonpharmacologicals to ease distressing symptoms, as well as thoughtful communication to support important elements of decision-making (e.g., autonomy, capacity determination, and surrogacy). ICU clinicians also must anticipate and address the needs of surviving relatives and marshal resources to mitigate the impact of their loss. Structured approaches are available to guide practice [4].

4. A decision to allow a natural death does not by itself mean that comfort is the patient’s exclusive goal or that intensive care interventions should be withheld/withdrawn

Patients’ goals of care can be multidimensional, encompassing a desire to allow natural death (e.g., DNR/DNAR) along with a preference to pursue life-prolonging interventions. Without explicit discussion, a DNR directive should not be interpreted as extending to interventions other than cardiopulmonary resuscitation. Clear, sensitive, and nuanced discussion of the patient’s true values, goals, and preferences is essential.

5. Critical illness and intensive care have profound impact on families as well as patients

The impact of critical illness is deeply felt by patients’ families, who experience psychological and physical distress as well as practical and financial burdens [5]. This can manifest acutely as depression, fear, anxiety, fatigue, anorexia, and early posttraumatic stress symptoms. Persistence of symptoms after the ICU can result in

*Correspondence: jde2134@columbia.edu

¹ Division of Pediatric Critical Care, Columbia University College of Physician and Surgeons, 3959 Broadway, CHN 10-24, New York, NY 10032, USA

Full author information is available at the end of the article

Table 1 ICU palliative care educational resources and clinical tools found online

	Online courses/ education	Webinars	In-person workshops	Clinical tools/ guidelines
American Association of Colleges of Nursing - The End-of-Life Nursing Education Consortium (http://www.aacn.nche.edu/el nec)				
American Association of Critical-Care Nurses (http://www.aacn.org/wd/practice/content/palliative-and-end-of-life-care.pcms?menu=practice)				
Center to Advance Palliative Care (https://www.capc.org)				
Coalition for Compassionate Care of California (http://coalitionccc.org)				
Education in Palliative and End-of-life Care (http://www.epec.net)				
European Association for Palliative Care (http://www.eapcnet.eu/Themes/Education.aspx)				
The IPAL-ICU Project (https://www.capc.org/ipal/ipal-icu/)				
National Consensus Project for Quality Palliative Care (http://www.nationalconsensusproject.org/NCP_Clinical_Practice_Guidelines_3rd_Edition.pdf)				
PCNOW (http://www.mypcnow.org)				
VitalTalk (http://vitaltalk.org/courses)				

worsening health-related quality-of-life and “post-intensive care unit syndrome-family” in families of patients who survive and bereaved families [6]. Risk and protective factors for family morbidity and burden have been investigated; some are potentially modifiable [6].

6. Palliative care can support ICU clinicians who experience moral or psychological distress in the course of caring for patients

ICU clinicians are at risk for moral distress (e.g., from conscientious objections) and psychological distress (e.g., grief and burnout). Primary and specialist palliative care offer a paradigm to help understand, anticipate, and address these effects. Strategies include strong interprofessional collaboration, support for resilience, attention to and normalization of grief and other emotions, mindfulness and meditation training, and forums to share experiences and reflections.

7. Educational resources and clinical tools are available to support delivery of high-quality palliative care in the ICU

Examples of online resources and tools for knowledge and skills supporting primary ICU palliative care are found in Table 1. Other resources include online courses (e.g., VitalTalk <http://vitaltalk.org/courses>) and published protocols to improve communication [7], studies demonstrating the effect of communication interventions [8, 9], and

published guidelines [3]. Palliative care specialists also can contribute importantly in educating the ICU team to assess and respond to palliative needs at the primary level.

8. The ICU nurse plays a crucial role in palliative care including communication between the health care team and the patient/family

High-quality ICU palliative care depends on crucial contributions made by ICU nurses, who are typically patients’ closest and most constant providers. Among other key roles, these nurses can help open lines of interprofessional communication with families, while also providing clarity, reassurance, consistency, and continuity in ICU family meetings [8, 10, 11]. Nurses’ participation in such meetings is an indicator of ICU care quality [12].

9. The quality of ICU palliative care can be measured

Domains of high-quality ICU palliative care have been defined by ICU patients, families, clinicians, and expert consensus, and have been operationalized as process measures for monitoring, feedback, and improvement [12, 13]. Opportunities to improve performance on such measures exist in many ICUs [14]. The relationship between process measures and outcomes of importance to ICU patients and families requires further investigation.

10. High-quality ICU palliative care can improve ICU metrics and patient-oriented outcomes

ICU-based palliative care interventions have been shown to improve utilization (e.g., length of stay), the quality and content of clinician communication, and family psychological well-being, satisfaction, and decision-making [9, 15–17]. Ongoing research is expanding the range of such interventions and comparing efficiency, effectiveness, and scalability. It will also be important to investigate which outcomes are most valid and responsive, which patients/families are most likely to benefit, and which interventions might be accompanied by unintended adverse effects [18].

Palliative care complements and augments intensive care in addressing the needs of critically ill patients and their families. Palliative care is based on need, not prognosis, and is optimally provided together with all appropriate life-prolonging or restorative care. Palliative care delivered by ICU clinicians and specialists contributes positively to outcomes of importance to ICU patients, families, and clinicians. Going forward, increasing evidence of the challenges facing survivors of critical illness and their families opens new opportunities for palliative care to support optimal critical care practice.

Author details

¹ Division of Pediatric Critical Care, Columbia University College of Physician and Surgeons, 3959 Broadway, CHN 10-24, New York, NY 10032, USA. ² Critical Care Service, Memorial Sloan Kettering Cancer Center, Weill Cornell Medical College, New York, USA. ³ Palliative Medicine Service, Memorial Sloan Kettering Cancer Center, Weill Cornell Medical College, New York, USA.

Compliance with ethical standards

Conflicts of interest

The authors have no potential financial or other conflicts of interest to disclose.

Received: 12 July 2016 Accepted: 30 July 2016

Published online: 9 August 2016

References

- Nelson JE, Bassett R, Boss RD, Brasel KJ, Campbell ML, Cortez TB, Curtis JR, Lustbader DR, Mulkerin C, Puntillo KA, Ray DE, Weissman DE (2010) Models for structuring a clinical initiative to enhance palliative care in the intensive care unit: a report from the IPAL-ICU Project (Improving Palliative Care in the ICU). *Crit Care Med* 38:1765–1772
- Downar J, Delaney JW, Hawryluck L, Kenny L (2016) Guidelines for the withdrawal of life-sustaining measures. *Intensive Care Med* 42:1003–1017
- Truog RD, Campbell ML, Curtis JR, Haas CE, Luce JM, Rubenfeld GD, Rushton CH, Kaufman DC, American Academy of Critical Care Medicine (2008) Recommendations for end-of-life care in the intensive care unit: a consensus statement by the American College [corrected] of Critical Care Medicine. *Crit Care Med* 36:953–963
- Treece PD, Engelberg RA, Crowley L, Chan JD, Rubenfeld GD, Steinberg KP, Curtis JR (2004) Evaluation of a standardized order form for the withdrawal of life support in the intensive care unit. *Crit Care Med* 32:1141–1148
- Schmidt M, Azoulay E (2012) Having a loved one in the ICU: the forgotten family. *Curr Opin Crit Care* 18:540–547
- Davidson JE, Jones C, Bienvenu OJ (2012) Family response to critical illness: postintensive care syndrome-family. *Crit Care Med* 40:618–624
- Curtis JR, White DB (2008) Practical guidance for evidence-based ICU family conferences. *Chest* 134:835–843
- Krimshtein NS, Luhrs CA, Puntillo KA, Cortez TB, Livote EE, Penrod JD, Nelson JE (2011) Training nurses for interdisciplinary communication with families in the intensive care unit: an intervention. *J Palliat Med* 14:1325–1332
- Sullivan AM, Rock LK, Gadmer NM, Norwich DE, Schwartzstein RM (2016) The impact of resident training on communication with families in the intensive care unit. Resident and family outcomes. *Ann Am Thorac Soc* 13:512–521
- Garrouste-Orgeas M, Max A, Lerin T, Grégoire C, Ruckly S, Kloeckner M, Brochon S, Pichot E, Simons C, El-Mhadri M, Bruel C, Philippart F, Fournier J, Tiercelet K, Timsit JF, Misset B (2016) Impact of proactive nurse participation in ICU family conferences: a mixed-method study. *Crit Care Med* 44:1116–1128
- Nelson JE, Cortez TB, Curtis JR, Lustbader, Mosenthal AC, Mulkerin C, DE Ray, Bassett R, Boss RD, Brasel KJ, Campbell ML, DE Weissman, Puntillo KA, The IPAL-ICU Project™ (2011) Integrating palliative care in the ICU: the nurse in a leading role. *J Hosp Palliat Nurs* 13:89–94
- Nelson JE, Mulkerin CM, Adams LL, Pronovost PJ (2006) Improving comfort and communication in the ICU: a practical new tool for palliative care performance measurement and feedback. *Qual Saf Health Care* 15:264–271
- Nelson JE, Puntillo KA, Pronovost PJ, Walker AS, McAdam JL, Ilaoa D, Penrod J (2010) In their own words: patients and families define high-quality palliative care in the intensive care unit. *Crit Care Med* 38:808–818
- Penrod JD, Pronovost PJ, Livote EE, Puntillo KA, Walker AS, Wallenstein S, Mercado AF, Swoboda SM, Ilaoa D, Thompson DA, Nelson JE (2012) Meeting standards of high-quality intensive care unit palliative care: clinical performance and predictors. *Crit Care Med* 40:1105–1112
- Aslakson R, Cheng J, Vollenweider D, Galusca D, Smith TJ, Pronovost PJ (2014) Evidence-based palliative care in the intensive care unit: a systematic review of interventions. *J Palliat Med* 17:219–235
- Lautrette A, Darmon M, Megarbane B, Joly LM, Chevret S, Adrie C, Barnoud D, Bleichner G, Bruel C, Choukroun G, Curtis JR, Fieux F, Galliot R, Garrouste-Orgeas M, Georges H, Goldgran-Toledano D, Jourdain M, Loubert G, Reignier J, Saida F, Souweine B, Vincent F, Barnes NK, Pochard F, Schlemmer B, Azoulay E (2007) A communication strategy and brochure for relatives of patients dying in the ICU. *N Engl J Med* 356:469–478
- Khandelwal N, Kross EK, Engelberg RA, Coe NB, Long AC, Curtis JR (2015) Estimating the effect of palliative care interventions and advance care planning on ICU utilization: a systematic review. *Crit Care Med* 43:1102–1111
- Carson SS, Cox CE, Wallenstein S, Hanson LC, Danis M, Tulsy JA, Chai E, Nelson JE (2016) Effect of palliative care-led meetings for families of patients with chronic critical illness: a randomized clinical trial. *JAMA* 316:51–62